

兔抗 WDR83 多克隆抗体

中文名称：兔抗 WDR83 多克隆抗体

英文名称： Anti-WDR83 rabbit polyclonal antibody

别名： MORG1

相关类别： 一抗

储存： 冷冻（-20℃）

抗原： WDR83

宿主： Rabbit

反应种属： Human, Mouse, Rat

标记物： Unconjugate

克隆类型： rabbit polyclonal

技术规格

| | |
|--------------------------|--|
| Background: | This gene encodes a member of the WD-40 protein family. The protein is proposed to function as a molecular scaffold for various multimeric protein complexes. The protein associates with several components of the extracellular signal-regulated kinase (ERK) pathway, and promotes ERK activity in response to serum or other signals. The protein also interacts with egl nine homolog 3 (EGLN3, also known as PHD3) and regulates expression of hypoxia-inducible factor 1, and has been purified as part of the spliceosome. Alternative splicing results in multiple transcript variants. |
| Applications: | ELISA, WB, IHC |
| Name of antibody: | WDR83 |

| | |
|------------------------------------|---|
| Immunogen: | Fusion protein of human WDR83 |
| Full name: | WD repeat domain 83 |
| Synonyms : | MORG1 |
| SwissProt: | Q9BRX9 |
| ELISA Recommended dilution: | 2000-5000 |
| IHC positive control: | Human thyroid cancer and human colon cancer |
| IHC Recommend dilution: | 50-200 |
| WB Predicted band size: | 34 kDa |
| WB Positive control: | 231 and A172 cells |
| WB Recommended dilution: | 200-1000 |



